

AD-A286 345



TATION PAGE

Form Approved
OMB No. 0704-0188

1. TO: (Agency or person to whom the report is submitted, including the name of the reviewing organization, searching existing data sources, reviewing the collection of information, and comments regarding this report estimate or any other aspect of this report.)
2. FROM: (Agency or person from whom the report is submitted, including the name of the reviewing organization, searching existing data sources, reviewing the collection of information, and comments regarding this report estimate or any other aspect of this report.)
3. MANAGEMENT AND BUDGET: (Agency or person from whom the report is submitted, including the name of the reviewing organization, searching existing data sources, reviewing the collection of information, and comments regarding this report estimate or any other aspect of this report.)

DATE
October 18, 19943. REPORT TYPE AND DATES COVERED
Technical Report

4. TITLE AND SUBTITLE

Synthesis and Structure of a Novel Bis-Stibole

6. AUTHOR(S)

Donald P. Hsu, Benjamin P. Warner, Richard A. Fisher, William M. Davis, and Stephen L. Buchwald

5. FUNDING NUMBERS

Grant
#N00014-91-J-1043
R&T Code 3135014
Harold E. Guard,
Program Officer

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Massachusetts Institute of Technology
Department of Chemistry
Cambridge, MA 02139

94-35063



9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Office of Naval Research
Department of the Navy
Chemistry Division, ONR 331
800 North Quincy Street
Arlington, VA 22217-566010. SPONSORING / MONITORING
AGENCY REPORT NUMBER

Technical Report No. 6

11. SUPPLEMENTARY NOTES

Prepared for Publication in Organometallics


12a. DISTRIBUTION / AVAILABILITY STATEMENT

Reproduction in whole or in part is permitted for any purpose of the United States Government. Approved for public release and sale, distribution unlimited.

12b. DISTRIBUTION CODE

13. ABSTRACT (Maximum 200 words)

The novel benzobis(stibole) **2a**, synthesized from bis(zirconacycle) **1**, crystallizes to place all three ring substituents in a *syn* orientation. Complex **2a** stacks in such a manner so that the central phenyl ring of two adjacent Sb_2C_{10} ring systems are nearly perfectly eclipsed.



94 11 14 012

14. SUBJECT TERMS

stibole, organo-antimony, organo-main group

15. NUMBER OF PAGES
10

16. PRICE CODE

17. SECURITY CLASSIFICATION
OF REPORT

Unclassified

18. SECURITY CLASSIFICATION
OF THIS PAGE

Unclassified

19. SECURITY CLASSIFICATION
OF ABSTRACT

Unclassified

20. LIMITATION OF ABSTRACT

UL

NSN 7540-01-280-5500

DTIC QUALITY INSURED

Standard Form 298 (Rev. 2/89)
Prescribed by ANSI Std. Z39-18
298-102

OFFICE OF NAVAL RESEARCH

GRANT #N00014-91-J-1043

**R&T Code 3135014
Harold E. Guard, Program Officer
Chemistry Division**

Technical Report No 6

**Synthesis and Structure
of a Novel Bis-Stibole**

by

**Donald P. Hsu, Benjamin P. Warner, Richard A. Fisher,
William M. Davis and Stephen L. Buchwald**

Prepared for Publication

in

Organometallics

**Massachusetts Institute of Technology
Department of Chemistry
Cambridge, MA 02139**

October 18, 1994

**Reproduction in whole or in part is permitted for any purpose of the United States
Government**

Synthesis and Structure of a Novel Bis-Stibole

Donald P. Hsu, Benjamin P. Warner, Richard A. Fisher, William M. Davis and Stephen L. Buchwald*

Massachusetts Institute of Technology

Cambridge, MA 02139 (USA)

The novel benzobis(stibole) **2a**, synthesized from bis(zirconacycle) **1**, crystallizes to place all three ring substituents in a syn orientation. Complex **2a** stacks in such a manner so that the central phenyl ring of two adjacent Sb_2C_{10} ring systems are nearly perfectly eclipsed.


Accession Number		1
Date		11/11/91
Author		Donald P. Hsu, Benjamin P. Warner, Richard A. Fisher, William M. Davis and Stephen L. Buchwald
Title		Synthesis and Structure of a Novel Bis-Stibole
Abstract		
Keywords		
Availability Codes		
Dist	Availability Codes	Special
A-1		

Revised

For several years we have been interested in the use of zirconacycles for the synthesis of novel main group derivatives.¹ Our efforts have focused principally on developing synthetic routes to cyclic compounds of the heavier main group elements for which there are few reliable methods.² We have been particularly interested in synthetic routes to stiboles, certain derivatives of which are thermochromic.³ In this account, we report the synthesis and solid state structure of some unusual bis-stibolobenzene derivatives.

We have previously reported the synthesis, characterization, and reactivity of zirconocene complexes of certain benzdiynes and these studies provide the starting point for the current investigation. We felt that bis-zirconacycle **1**, prepared as previously described,⁴ would provide us with a ready entry for the synthesis of unique bis-main group heterocycles by use of the procedure developed by Fagan and Nugent.⁵ Treatment of toluene solutions of **1** with two equivalents phenylantimony dichloride, gives the isomeric phenyl stiboles **2** (Scheme 1).⁶ Filtration of the reaction mixture through alumina (III) provides **2** in 71% yield as a mixture of *syn* (**2a**) and *anti* (**2b**) isomers (**2a/2b** = 55/45). Isomer **2a** can be preferentially crystallized (rectangular

Scheme 1



prisms) from cold hexane/toluene solutions of the mixture of isomeric metallacycles. Recrystallization of **2** from ether generates two distinct crystal types, rectangular prisms of **2a** and small needles of **2b**. The crystal types may be physically separated to give small amounts of each isomer. Once purified, solutions of the isomers are stable with respect to interconversion at temperatures up to 150 °C, at which point the compounds

begin to decompose. However, **2** is not configurationally stable if exposed to Lewis acids.⁷ When pure **2a** was adsorbed onto silica gel for 1 h, then eluted with ether, the recovered compound **2** was found to be a mixture of **2a:2b** in a 82:18 ratio.

The solid state structure of **2a** was determined by a single crystal x-ray diffraction study.⁸ The ORTEP and packing diagrams are shown in Figures 1 and 2. The bond distances and angles within the ring system are in the range found in stiboles which have been previously reported.^{1,3a-c} The Sb₂C₁₀ ring system is virtually planar. The bond angles about the antimony atom are narrow as might be expected from consideration of the hybridization of the antimony center in stibines.^{3a} Quite unexpectedly however, the methyl group of the ring methoxy substituent is oriented *syn*

Figure 1. Molecular structure of compound **2a** with partial listing of bond distances and angles.

to the two phenyl groups located on the antimony centers. The likely reason for this structural feature is apparent from the examination of the packing diagram (Figure 2)

Figure 2. Packing diagram of compound **2a**

and consideration of the close intermolecular contacts. Two Sb₂C₁₀ ring systems lie parallel to each other, 3.6 Å apart. This proximity forces the methoxy carbon atoms to reside away from neighboring Sb₂C₁₀ ring system and toward the phenyl substituents. The interactions between the two Sb₂C₁₀ rings are shown in Figure 3 in which the phenyl rings have been removed for clarity. Of particular interest is that the central six membered rings of the two molecules, shown in Figure 3, are nearly perfectly eclipsed.

Figure 3. Orientation of the symmetry related Sb₂C₁₀ rings. The phenyl rings have been omitted.

This is in contrast to Jorgensen's recent theoretical results for the structure of the benzene-anthracene complex.⁹ This interesting difference may be due to crystal packing forces. The corresponding stibacyclopentadiene units are approximately anti with Sb₁ and Sb₂ lying between C1 and C2 and C7 and C8 of the opposite ring respectively. This orientation minimizes the unfavorable coulombic interactions between the lone pairs of related antimony centers of the two units.

In summary, we have prepared the first example, to our knowledge, of bis-metallacyclopentadienes of a main group element. Investigations are underway in our laboratory to further explore the use of zirconacyclic intermediates as vehicles for the preparation of new and interesting main group compounds.

Experimental Section

Diphenyldistibolobenzenes: (2a, 2b):

Two dry Schlenk flasks were charged with **1** (0.400 g, 0.61 mmol) and phenylantimony dichloride (0.329 g, 1.22 mmol) separately, in a nitrogen-filled glove box. The flasks were removed from the glove box and attached to a Schlenk line under argon, and toluene (10 mL) was added to each. The solution of **1** was added to the phenylantimony dichloride solution *via* cannula at room temperature. The color of the reaction mixture quickly changed from red to pale yellow. The reaction mixture was stirred for 2 h, then a sample was removed by pipette. At this point, the reaction was judged to be complete (¹H NMR), with the product being a 55:45 ratio of **2a**:**2b**. The reaction solution was then filtered through a plug of alumina (III) with an additional 50 mL of toluene used to wash the material through the column. The solvent was removed *in vacuo* to leave a yellow solid (0.265 g, 0.434 mmol, 71%). This material was dissolved in approximately 0.5 mL of toluene and 2.5 mL of hexane (with gentle heating to 50 °C to dissolve), and allowed to crystallize, under argon, at -20 °C to yield several large, blocky, yellow crystals (0.062g, 0.102 mmol, 17%). Further attempts to separate

the diastereomers from the remaining material by fractional crystallization were unsuccessful. In a separate experiment, a mixture of **2a** and **2b** was crystallized from ether to yield two distinct forms of crystals: rectangular (**2a**) and needle-type crystals (**2b**). A small amount of the each isomer (**2b**) was physically separated and used for full characterization: **2a**: ^1H NMR (300 MHz, C_6D_6) δ 7.5-7.4 (m, 4H), 7.32 (s, 1 H), 7.0-6.9 (m, 6 H), 3.45 (s, 3 H), 2.05 (s, 6 H) and 2.03 (s, 6 H); ^{13}C NMR (75 MHz, CDCl_3) δ 166.3, 158.3, 147.1, 145.6, 137.5, 135.6, 131.0, 128.7, 128.4, 116.8, 60.7, 18.6 and 14.8; IR (KBr) 3059, 2987, 2907, 2848, 1563, 1510, 1475, 1430, 1364, 1350, 1317, 1193, 1063, 1035, 1018, 998, 855, 730, 695 and 450 cm^{-1} ; Anal. Calcd. for $\text{C}_{27}\text{H}_{26}\text{OSb}_2$: C, 53.16; H, 4.3. Found: C, 53.05; H, 4.36. **2b**: ^1H NMR (300 MHz, CDCl_3) δ 7.46-7.42 (m, 4 H), 7.26-7.18 (m, >6 H solvent included), 3.61 (s, 3 H), 2.21 (s, 6 H) and 2.19 (s, 6 H); ^{13}C NMR (75 MHz, CDCl_3) δ 166.3, 158.3, 147.1, 145.6, 137.3, 135.6, 130.7, 128.8, 128.5, 116.8, 60.5, 18.5 and 14.8; IR (KBr) 3060, 2986, 2929, 2908, 2842, 1565, 1512, 1476, 1430, 1354, 1318, 1063, 1036, 1019, 998, 855, 729, 695 and 450 cm^{-1} ; Anal. Calcd. for $\text{C}_{27}\text{H}_{26}\text{OSb}_2$: C, 53.16; H, 4.3. Found: C, 53.37; H, 4.36.

Acknowledgment. This work was supported, in part, by the Office of Naval Research, and by Union Carbide (Innovation Recognition Award), whom we gratefully acknowledge. SLB acknowledges additional support as a Fellow of the Alfred P. Sloan Foundation (1988-92) and a Camille & Henry Dreyfus Teacher-Scholar (1989-94).

Supplementary Material Available: ORTEP diagram of **2a**, tables of bond distances and angles for **2a**, and a table of final positional and thermal parameters for **2** (29 pages). Ordering information is given on any current masthead page.

References and Notes:

1. (a) Buchwald, S.L.; Fisher, R.A.; Davis, W.M. *Organometallics* **1989**, *8*, 2082.
 (b) Buchwald, S.L.; Fisher, R.A.; Foxman, B.A. *Angew.Chem. Int. Ed. Eng.* **1990**, *29*, 771. (c) Fisher, R.A.; Nielsen, R.B.; Davis, W.M.; Buchwald, S.L. *J. Am. Chem. Soc.* **1991**, *113*, 165. (d) Spence, R.E.v.H.; Hsu, D.P.; Davis, W.M.; Buchwald, S.L.; Richardson, J.F. *Organometallics* **1992**, *11*, 3492.

2. (a) Aylett, B.J. *Organometallic Compounds, Fourth Edition, Part Two*, Wiley: New York, **1979**. (b) Duak, G.O.; Freedman, L.D. *Organometallic Compounds of Arsenic, Antimony, and Bismuth*. Wiley: New York, **1970**. (c) Atkinson, R.E. in *Comprehensive Heterocyclic Chemistry*, (Ed: O. Meth-Cohn) Pergamon: Oxford, **1984**; Vol. 1, pp 539-560. (d) Annitage, D.A. in *Comprehensive Heterocyclic Chemistry*, (Ed: O. Meth-Cohn) Pergamon: Oxford, **1984**; Vol. 1, pp 614-617.
 (d) Markl, G.; Hauptman, H.; Merz, A. *J. Organomet. Chem.* **1983**, *249*, 335.

3. (a) Ashe, A.J. III; Butler, W.; Diephouse, T.R. *J. Am. Chem. Soc.* **1981**, *103*, 207.
 (b) Ashe, A.J. III; Kausch, C.M.; Eisenstein, O. *Organometallics* **1987**, *6*, 1185.
 (c) Bel'skii, V.K. *Zh. Strukt. Khim.* **1984**, *25*, 138. (d) Ashe, A.J. III *Adv. Organometal. Chem.* **1990**, *30*, 77.

4. Hsu, D.P.; Lucas, E.A.; Buchwald, S.L. *Tetrahedron Lett.* **1990**, *31*, 5563. Hsu, D.P., PhD Thesis, Department of Chemistry, MIT, 1991.

5. (a) Fagan, P.J.; Nugent, W.A. *J. Am. Chem. Soc.* **1988**, *110*, 2310. (b) Fagan, P.J.; Nugent, W.A.; Calabrese, J.C. *J. Am. Chem. Soc.* **1994**, *116*, 1880. (c) Fagan, P.J.; Burns, E.G.; Calabrese, J.C. *J. Am. Chem. Soc.* **1988**, *110*, 2979. See also (d) Fryzuk, M.D.; Bates, G.S.; Stone, C. *Tetrahedron Lett.* **1986**, *27*, 1537. (e) Fryzuk, M.D.; Bates, G.S.; Stone, C. *J. Org. Chem.* **1987**, *53*, 2334.

- (f) Fryzuk, M.D.; Bates, G.S.; Stone, C. *J. Org. Chem.* **1988**, *53*, 4425. (g) Doxsee, K.M.; Shen, G.S.; Knobler, C.B. *J. Am. Chem. Soc.* **1989**, *111*, 9129. (h) Tumas, W.; Suriano, J.A.; Harlow, R.L. *Angew. Chem. Int. Ed. Engl.* **1990**, *29*, 75. (i) Cole, T.E.; Quintanilla, R.; Rodewald, S. *Organometallics* **1991**, *10*, 3777. (j) Boutonnet, F.; Dufour, N.; Straw, T.; Igau, A.; Majoral, J.P. *Organometallics* **1991**, *10*, 3777. and references 1a-c.
6. When **1** was treated with 2 equiv. SbCl₃ or SbBr₃, the analogous chloro- and bromostiboles were formed as quite insoluble powders, which we were unable to isolate in sufficient purity.
7. (a) Campbell, I.G.M. *J. Chem. Soc.* **1947**, 4. (b) Campbell, I.G.M. *J. Chem. Soc.* **1952**, 4448. (c) Campbell, I.G.M.; Morrill, D.J. *J. Chem. Soc.* **1955**, 1662.
8. Crystals of **2a** are monoclinic; space group P2₁/n (#14); a = 12.284(2), b = 14.281(3), c = 13.898(3) Å; β = 95.31(20°; Z = 4; ρ_{calc} = 1.669 g/cm³. A total of 5790 unique reflections were collected using a Rigaku AFC6R diffractometer with graphite monochromated Mo Kα radiation and a 12KW rotating anode generator. The structure was solved using a combination of the Patterson method and direct methods. The final cycle of full-matrix least-squares refinement was based on 2832 observed reflections for which I > 3.00σ(I) and 271 variable parameters and converged with R = 0.037 and R_w = 0.053. Anisotropic thermal parameters were assigned to non-hydrogen atoms while hydrogen atoms were included as fixed contributions to the structure factors. Further details of the crystal structure investigation are available on request from the Director of the Cambridge Crystallographic Data Centre, University Chemical Laboratory, Lensfield Road, GB-Cambridge CB2 1EW (UK), on quoting the full journal citation.

9. Jorgenson, W.L.; Severance, D.L. *J. Am. Chem. Soc.* 1990, 112, 4768.

TECHNICAL REPORT DISTRIBUTION LIST - GENERAL

Office of Naval Research (1)*
Chemistry Division, ONR 331
800 North Quincy Street
Arlington, Virginia 22217-5660

Dr. Richard W. Drisko (1)
Naval Facilities & Engineering
Service Center
Code L52
Port Hueneme, CA 93043

Defense Technical Information
Center (2)
Building 5, Cameron Station
Alexandria, VA 22314

Dr. Eugene C. Fischer (1)
Code 2840
Naval Surface Warfare Center
Carderock Division Detachment
Annapolis, MD 21402-1198

Dr. James S. Murday (1)
Chemistry Division, Code 6100
Naval Research Laboratory
Washington, D.C. 20375-5320

Dr. Bernard E. Douda (1)
Crane Division
Naval Surface Warfare Center
Crane, Indiana 47522-5000

Dr. John Fischer, Director (1)
Chemistry Division, C0235
Naval Air Weapons Center
Weapons Division
China Lake, CA 93555-6001

Dr. Peter Seligman (1)
Naval Command, Control and
Ocean Surveillance Center
RDT&E Division
San Diego, CA 92152-5000

* Number of copies to forward